



LET'S GET STARTED!

A step-by-step guide to help you prepare for your first dose of PONVORY®

INDICATION

WHAT IS PONVORY®?

PONVORY® is a prescription medicine that is used to treat relapsing forms of multiple sclerosis, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

It is not known if PONVORY® is safe and effective in children.

Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).

We Know You've Got a Lot to Juggle, So Let Us Help. We Created This Kit for You to Navigate Your First 14 Days on PONVORY®

What's inside?

✓ **A Guide to Starting PONVORY®**

Understand the importance of taking the tablets in your Starter Pack exactly as directed

✓ **Starter Pack Instructions and Diagram**

Learn how to stay on track with your 14-Day Starter Pack

IMPORTANT SAFETY INFORMATION

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT PONVORY®?

PONVORY® may cause serious side effects, including:

- **Infections** – PONVORY® can increase your risk of serious infections that can be life-threatening and cause death. PONVORY® lowers the number of white blood cells (lymphocytes) in your blood. This will usually go back to normal within 1 to 2 weeks of stopping treatment. Your healthcare provider should review a recent blood test of your white blood cells before you start taking PONVORY®.

DO YOU PREFER THE WEB?

Activate this QR code using your smartphone and view online information about starting PONVORY®.

Open the camera on your phone

Point your camera at the QR code

Follow the prompt that appears on-screen to visit ponvoryus.com



Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).





Not an actual patient

Part 1: Getting Started

How to Use the PONVORY®
14-Day Starter Pack 5

Part 2: Staying on Track

A Diagram to Help You Stay
on Track During Your First
14 Days on PONVORY® 12

IMPORTANT SAFETY INFORMATION (continued)

Call your healthcare provider right away if you have any of these symptoms of an infection during treatment and for 1 to 2 weeks after your last dose of PONVORY®:

- fever
- tiredness
- body aches
- chills
- nausea
- vomiting
- headache with fever, neck stiffness, sensitivity to light, nausea, or confusion (these may be symptoms of meningitis, an infection of the lining around your brain and spine)

Your healthcare provider may delay starting or may stop your PONVORY® treatment if you have an infection.

Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).

 **Ponvory**
(ponesimod) once-daily tablets



Not an actual patient

Part 1

Getting Started

It's important to follow the dosing instructions for PONVORY®, which is why we've included a step-by-step guide to help you stay on track and avoid missing a dose. **Be sure to write down the day and date you start taking PONVORY®.**

IMPORTANT SAFETY INFORMATION (continued)

- **Slow heart rate (bradycardia or bradyarrhythmia) when you start taking PONVORY®.** PONVORY® can cause your heart rate to slow down, especially after you take your first dose. You should have a test to check the electrical activity of your heart called an electrocardiogram (ECG) before you take your first dose.

Please see full Important Safety Information throughout this brochure. [Click here for full Prescribing Information](#) including [Medication Guide](#).

 **Ponvory**
(ponesimod) once-daily tablets

Getting Started



The PONVORY® 14-Day Starter Pack slowly increases your dose of PONVORY® so your body can adjust to the new medicine.

2 mg → 10 mg

Your Starter Pack is clearly labelled with the day you will take each pill along with the pill's dose. For example, on Day 1 your dose will be 2 mg and by Day 14 your dose will be 10 mg.



per day

You will take only 1 pill at the same time each day when using the Starter Pack, with or without food. **Make sure that you're on the right day so you take your pills in order.**



After you complete the Starter Pack, you will start taking the recommended dose of 20 mg of PONVORY®. This is also called your maintenance dose. You will receive a separate prescription bottle with your 20 mg maintenance dose.

Depending on your pretests and/or cardiac history, you might also be asked to complete **first dose monitoring**. Your healthcare professional will inform you if this monitoring is required.



Not actual patients

IMPORTANT SAFETY INFORMATION (continued)

Only Start your treatment with PONVORY® using the Starter Pack. You must use the PONVORY® Starter Pack by slowly increasing the dose over a 14-day period to help reduce the effect of slowing of your heart rate. It is important to follow the recommended dosing instructions.

Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).

**Ponvory**
(ponesimod) once-daily tablets



Not an actual patient

Managing Your Dose

- Try your best not to miss a dose of PONVORY®.
- If you miss taking 1, 2, or 3 PONVORY® pills in a row in the 14-day Starter Pack, continue treatment by taking the **first** dose you missed as soon as you remember. Then, take 1 pill a day to continue with the Starter Pack dose as planned.
- If you miss taking 1, 2, or 3 PONVORY® pills in a row while taking the 20 mg maintenance dose, continue treatment with the 20 mg maintenance dose.
- If you miss taking PONVORY® for 4 or more days in a row while taking your Starter Pack or maintenance dose, you need to restart treatment with a new 14-day Starter Pack. You should call your healthcare professional right away.

Do not change your dose or stop taking PONVORY® unless your healthcare professional tells you to.

IMPORTANT SAFETY INFORMATION (continued)

Call your healthcare provider if you experience the following symptoms of slow heart rate:

- dizziness
- lightheadedness
- feeling like your heart is beating slowly or skipping beats
- shortness of breath
- confusion
- chest pain
- tiredness

Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).

 **Ponvory**
(ponesimod) once-daily tablets

Part 2

Staying on Track

Staying on top of taking PONVORY® may help you do more of what matters most.

IMPORTANT SAFETY INFORMATION (continued)

Do not take PONVORY® if you:

- have had a heart attack, chest pain called unstable angina, stroke or mini-stroke (transient ischemic attack or TIA), or certain types of heart failure in the last 6 months.
 - have certain types of heart block or irregular or abnormal heartbeat (arrhythmia) unless you have a pacemaker.
- Talk to your healthcare provider if you have any of these conditions, or do not know if you have any of these conditions.

A Diagram to Help You Stay on Track During Your First 14 Days on PONVORY®

As you go through your 14-Day Starter Pack, follow the instructions and arrows on your Starter Pack, which are reprinted with more details on the back of this card to make sure you're on track.

Tips to help you get started:


- Write down the day of the week and date you start taking your Starter Pack
- Set a daily alarm to remind you to take your medication and start to build a habit
- Remember your motivation for treating your multiple sclerosis (MS) and how treatment can help you reach your goals

Please see full Important Safety Information throughout this brochure. [Click here for full Prescribing Information](#) including [Medication Guide](#).



Using Your Starter Pack

This is new for you, so we want to help make sure you're taking the right dose on the right day.



This diagram is based off of an image of your Starter Pack.

*We've added **red arrows** to help you understand the order you should take the tablets in your Starter Pack.*

- **Take 1 pill each day**
- **Your start date is Day 1.** Take the pill labeled **Day 1**
- Continue with **Days 2-6** following the arrows closely between **Days 2-3** and **Days 4-5**
- Follow the arrow closely to get from **Day 6** to **Day 7**
- For **Days 7-9**, follow the horizontal row along the bottom
- For **Days 10-14**, follow the arrows closely between each day

The tablets vary in color and size based on the dosage strength.

Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).

 **Ponvory**
(ponesimod) once-daily tablets

Important Safety Information (continued)

Before you take PONVORY®, tell your healthcare provider about all your medical conditions, including if you:

- have a fever or infection, or you are unable to fight infections due to a disease or taking medicines that lower your immune system.
- have had chicken pox or have received the vaccine for chicken pox. Your healthcare provider may do a blood test for the chicken pox virus. You may need to get the full course of vaccine for chicken pox and then wait 1 month before you start taking PONVORY®.
- have slow heart rate.
- have an irregular or abnormal heartbeat (arrhythmia).
- have a history of stroke.
- have heart problems, including a heart attack or chest pain.
- have high blood pressure.
- have breathing problems, including during your sleep (sleep apnea).
- have liver problems.
- had or now have a type of skin cancer called basal cell carcinoma (BCC), melanoma, or squamous cell carcinoma
- have eye problems, especially an inflammation of the eye called uveitis.
- have diabetes.
- are pregnant or plan to become pregnant. PONVORY® may harm your unborn baby. Talk with your healthcare provider if you are pregnant or plan to become pregnant. If you are a woman who can become pregnant, you should use effective birth control during your treatment with PONVORY® and for 1 week after you stop taking PONVORY®. Talk to your healthcare provider about what method of birth control is right for you during this time. Tell your healthcare provider right away if you do become pregnant while taking PONVORY® or within 1 week after you stop taking PONVORY®.
- are breastfeeding or plan to breastfeed. It is not known if PONVORY® passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take PONVORY®.

Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements.

Using PONVORY® and other medicines together may affect each other causing serious side effects. Especially tell your healthcare provider if you take or have taken: Medicines to control your heart rhythm (antiarrhythmics), or blood pressure (antihypertensives), or heart-beat (such as calcium channel blockers or beta-blockers); medicines that affect your immune system, such as alemtuzumab; and medicines such as rifampin, phenytoin, or carbamazepine.

You should not receive **live** vaccines during treatment with PONVORY®, for at least 1 month before taking PONVORY®, and for 1 to 2 weeks after you stop taking PONVORY®. If you receive a live vaccine, you may get the infection the vaccine was meant to prevent. Vaccines may not work as well when given during treatment with PONVORY®.

Talk with your healthcare provider if you are not sure if you take any of these medicines.

HOW SHOULD I TAKE PONVORY®?

- Take PONVORY® exactly as your healthcare provider tells you to take it.
- Take PONVORY® 1 time each day.
- Swallow PONVORY® tablets whole.
- Take PONVORY® with or without food.
- Do not stop taking PONVORY® without talking with your healthcare provider first.
- Do not skip a dose.
- Start taking PONVORY® with a 14-day starter pack.
- If you miss taking 1, 2, or 3 tablets in a row of PONVORY® in the 14-day starter pack, continue treatment by taking the first dose you missed. Take 1 tablet as soon as you remember. Then, take 1 tablet a day to continue with the starter pack dose as planned.
- If you miss taking 1, 2, or 3 tablets in a row of PONVORY® while taking the 20 mg maintenance dose, continue treatment with the 20 mg maintenance dose.
- If you miss taking 4 or more tablets in a row of PONVORY®, while taking the 14-day starter pack or the 20 mg maintenance dose, you need to restart treatment with a new 14-day starter pack. Call your healthcare provider if you miss 4 or more doses of PONVORY®. Do not restart PONVORY® after stopping it for 4 or more days in a row without talking to your healthcare provider. If you have certain heart conditions, you may need to be monitored by your healthcare provider for at least 4 hours when you take your next dose.

Please see full Important Safety Information throughout this brochure. [Click here for full Prescribing Information including Medication Guide.](#)

 **Ponvory**
(ponesimod) once-daily tablets

Important Safety Information (continued)

What are the possible side effects of PONVORY®?

PONVORY® may cause serious side effects, including:

- **breathing problems.** Some people who take PONVORY® have shortness of breath. Call your healthcare provider right away if you have new or worsening breathing problems.
- **liver problems.** PONVORY® may cause liver problems. Your healthcare provider should do blood tests to check your liver before you start taking PONVORY®. Call your healthcare provider right away if you have any of the following symptoms of liver problems:
 - unexplained nausea
 - vomiting
 - stomach (abdominal) pain
 - tiredness
 - loss of appetite
 - yellowing of the whites of your eyes or skin
 - dark urine
- **increased blood pressure.** Your healthcare provider should check your blood pressure during treatment.
- **types of skin cancer called basal cell carcinoma (BCC), melanoma, and squamous cell carcinoma.** Certain types of skin cancer have happened with drugs in the same class. Tell your healthcare provider if you have any changes in the appearance of your skin, including changes in a mole, a new darkened area on your skin, a sore that does not heal, or growths on your skin, such as a bump that may be shiny, pearly white, skin-colored, or pink. Your doctor should check your skin for any changes during treatment with PONVORY®. Limit the amount of time you spend in sunlight and ultraviolet (UV) light. Wear protective clothing and use a sunscreen with a high sun protection factor.
- **a problem with your vision called macular edema.** Tell your healthcare provider about any changes in your vision. Your healthcare provider should test your vision before you start taking PONVORY® and any time you notice vision changes during treatment with PONVORY®. Your risk of macular edema is higher if you have diabetes or have had an inflammation of your eye called uveitis. Call your healthcare provider right away if you have any of the following symptoms:
 - blurriness or shadows in the center of your vision
 - a blind spot in the center of your vision
 - sensitivity to light
 - unusually colored (tinted) vision

- **swelling and narrowing of the blood vessels in your brain.** A condition called Posterior Reversible Encephalopathy Syndrome (PRES) has happened with drugs in the same class. Symptoms of PRES usually get better when you stop taking PONVORY®. However, if left untreated, it may lead to a stroke. Call your healthcare provider right away if you have any of the following symptoms:

- sudden severe headache
- sudden confusion
- sudden loss of vision or other changes in vision
- seizure

- **severe worsening of multiple sclerosis (MS) after stopping PONVORY®.** When PONVORY® is stopped, symptoms of MS may return and become worse compared to before or during treatment. Always talk to your healthcare provider before you stop taking PONVORY® for any reason. Tell your healthcare provider if you have worsening symptoms of MS after stopping PONVORY®.

The most common side effects of PONVORY® include:

- upper respiratory tract infections
- elevated liver enzymes (abnormal liver tests)
- high blood pressure (hypertension)

These are not all the possible side effects of PONVORY®. For more information, ask your healthcare provider or pharmacist. See “What is the most important information I should know about PONVORY®?”

Tell your doctor if you have any side effect that bothers you or that does not go away.

Call your doctor for medical advice about side effects. You are also encouraged to report side effects to the FDA: visit <http://www.fda.gov/medwatch> or call 1-800-FDA-1088. You may also report side effects to Vanda Pharmaceuticals, Inc., at 1-833-933-9331.

Please see full Important Safety Information throughout this brochure. Click here for full [Prescribing Information](#) including [Medication Guide](#).

Trademarks are those of their respective owners.

 **Ponvory**
(ponesimod) once-daily tablets

Not an actual patient



For more information,
visit ponvoryus.com

Please see full Important Safety Information throughout this brochure.
Click here for full [Prescribing Information](#) including [Medication Guide](#).